Final Examination Calculus I (MATH-156), Fall 2005

Max Points: 200

1. Draw the graph of the function with domain [0, 4] and having the following properties:

(i)
$$f(0) = -1$$
, $f(2) = 2$, and $f(4) = 1$.

(ii)
$$\lim_{x\to 1^-} f(x) = 1$$
.

(iii)
$$\lim_{x \to 1^+} f(x) = 3$$
.

(iv)
$$\lim_{x\to 4^-} f(x) = 0$$
.

2. (a) Prove that
$$\lim_{x\to -9} \left(\frac{x^2-81}{x+9}\right) = -18$$
.

(b) Suppose
$$\lim_{x\to\sqrt{2}} f(x) = 3$$
, and $\lim_{x\to\sqrt{2}} (fg)(x) = -\sqrt{2}$. show that $\lim_{x\to\sqrt{2}} g(x)$ exists and find $\lim_{x\to\sqrt{2}} g(x)$.

3. Show that

$$f(x) = \begin{cases} x^2 + 3, & x \le 1\\ x + 3, & x > 1 \end{cases}$$

is continuous, but not differentiable at x = 1. Sketch the graph of f.

4. Find the indicated derivative. Simplify your answers.

(i)
$$f(x) = \frac{(3-2x)^{4/3}}{x^2}$$
, find $f'(x)$.

(ii)
$$f(x) = \ln \frac{x^3}{\cos x}$$
, find $f'(x)$

(ii)
$$f(x) = \ln \frac{x^3}{\cos x}$$
, find $f'(x)$.
(iii) $f(x) = \pi^{(\sin x - \cos x)}$, find $f'(\frac{\pi}{4})$.

(iv)
$$f(x) = (\ln x)e^{3x}$$
, find $f'(1)$.

5. Evaluate the limit

(i)
$$\lim_{x\to 0} \frac{x^2 e^x}{\sin^2 3x}$$

(ii)
$$\lim_{x\to\infty} x \sin(\frac{\pi}{x})$$
.

6. (a) Use implicit differentiation to find $\frac{dy}{dx}$:

$$x^3 + y^3 = 3xy^2$$

(b) Assume that x and y are differentiable functions of t. Find $\frac{dy}{dt}$ in terms of x, y and $\frac{dx}{dt}$:

$$x \sin y = 2$$

7. (a) Find the absolute extrema values of

$$f(x) = (x-2)^2(x+3)^2$$
 on $(-\infty, \infty)$.

Compute also inflection points (if any), and provide a brief sketch of the curve.

- (b) Given the polynomial $f(x) = x^6 2x^2 + x$ on the interval [0, 2], show that the polynomial $6x^5 - 4x + 1$ has a zero on the open interval (0, 2).
- 8. A container with square base, vertical sides and open top is to be made from 1000 ft² of the material. Find the dimensions of the container with greatest volume.
- 9. Find
 - (a) $\int 3x^2(x^3+5)^9 dx$
 - (b) $\int \sin^4 x \cos x \, dx$
- 10. (a) Evaluate $\int_1^3 f(x) dx$, when

$$f(x) = \begin{cases} x & \text{for } 1 \le x \le 2\\ 2 & \text{for } 2 \le x \le 3 \end{cases}$$

(b) Let

$$g(x) = \int_0^x \frac{\cos t}{t^2 + 3} dt$$

Find (i) g'(0) and (ii) g''(0).